

Loss of the Delta's historical landscape as an ecological stressor

*Notes compiled by Brian Atwater with help from Elizabeth Canuel and Jeffrey Mount,
Delta Independent Science Board*

Landscape lost

At the time of the California Gold Rush, tidal wetlands upstream of Suisun Bay occupied about as much area as did the open water and mudflats of San Francisco, San Pablo, and Suisun Bays combined. Together with the Delta's channels, lakes, and natural levees, these Delta wetlands covered an estimated 1400 square kilometers^{1, 2}.

The Delta's vast marshy plains, served by networks of meandering tidal channels, adjoined gallery forests along the Sacramento River and its distributaries³, received outflow from riverine flood basins that held ponds and lakes^{4, 5}, surrounded sand mounds near Isleton and Brentwood⁶, abounded in willow-fern swamps along distributaries of the San Joaquin River⁷, and graded into the seasonally flooded toes of alluvial fans⁸. The structure of this bygone landscape is becoming clearer through examination of old maps, surveyors' notes, newspaper accounts, travel diaries, and photographs this bygone landscape⁸.

Some 20 square kilometers of Delta tidal wetland remained six decades ago⁹. Today, as they did then, the remnants dot dredge-cut waterways.

On assessing ecological stresses from this landscape loss

1. Assume that the near-total disappearance of the Delta's historical landscape abets other stressors, as in the "regime shift" hypothesis¹⁰.
2. Seek clues to ecological functions of the bygone landscape through study of the Delta's historical descriptions^{5, 9} and geological deposits¹¹⁻¹³.
3. Look for these functions in restorations that play the country-music song backwards:
 - a. Draw analogies with the Delta's new wetlands^{14, 15}. Levee failure at Liberty Island has yielded tidal mud flats, marshes, and dead-end sloughs believed beneficial to Delta smelt^{16, 17}.
 - b. Invert guiding principles of habitat restoration. According to guidelines proposed for the Puget Sound Nearshore Ecosystem Restoration Project¹⁸, restoration success is to likely to increase with the size and connectivity of the restored ecosystems, and with attempts to mimic their roles at landscape scale.
4. Make analogies with inferred effects of habitat loss at other estuaries. Diking of the Columbia River's freshwater wetlands may have harmed Chinook salmon by limiting export of vascular-plant detritus¹⁹.

References cited

1. Gilbert, G. K. Hydraulic-mining debris in the Sierra Nevada. U.S. Geological Survey Professional Paper 105, 154 p. (1917).
2. Atwater, B. F., Conrad, S. G., Dowden, J. N., Hedel, C. W., MacDonald, R. L., Savage, W. History, landforms, and vegetation of the estuary's tidal marshes, *in* San Francisco Bay: the urbanized estuary (ed Conomos, T. J.) 347-385, http://www.estuaryarchive.org/archive/conomos_1979/18/ (Pacific Division, American Association for the Advancement of Science, San Francisco, Calif., 1979).
3. Jepson, W. L. The riparian botany of the lower Sacramento. *Erythea* 1, 238-246 (1893).
4. Thompson, J. Settlement geography of the Sacramento - San Joaquin Delta. Ph.D. thesis, Stanford Univ., Stanford, Calif. 551 p. (1957).
5. Whipple, A., Grossinger, R., Rankin, D. & Collins, J. The historical Yolo Basin landscape. Oral Abstracts, Bay-Delta Science Conference, September 27-29, 2010, <http://www.baydeltascienceconf.com/sciconf2010/?q=node/414> (2010).
6. Atwater, B. F. Geologic maps of the Sacramento - San Joaquin Delta. 1:24,000, U.S. Geological Survey Miscellaneous Field Studies Map MF-1411 (1982).
7. Mason, H. L. Untitled report on floristics of the Sacramento - San Joaquin Delta. 84 p. On file at San Francisco Estuary Institute (undated).
8. Grossinger, R., Whipple, A., Rankin, D. & Collins, J. Historical Delta habitat mosaics: conceptual models for building a diverse and resilient future Delta. Oral Abstracts, Bay-Delta Science Conference, September 27-29, 2010, <http://www.baydeltascienceconf.com/sciconf2010/?q=node/414> (2010).
9. Cosby, S. W. Soil survey of the Sacramento - San Joaquin Delta. U.S. Department of Agriculture, Bureau of Chemistry and Soils series 1935, no. 41, 48 p. (1941).
10. Baxter, R., Breuer, R. Brown, L., Conrad, L., Feyrer, F., Fong, S., Gehrts, K., Grimaldo, L., Herbold, B., Hrodey, P., Mueller-Solger, A., Sommer, T., Souza, K. Interagency Ecological Program 2010 Pelagic Organism Decline work plan and synthesis of results. <http://www.water.ca.gov/iep/docs/FinalPOD2010Workplan12610.pdf> (2010).
11. Alpers, C. N., Drexler, J. Z., Neymark, L. A., Paces, J. B. & Taylor, H. E. Peat deposits of the Sacramento-San Joaquin Delta, California: an archive for anthropogenic contamination of mercury, lead, and other trace metals. *Eos* (Transactions, American Geophysical Union) 89, Abstract B13B-0441 (2008).
12. Drexler, J. Z., Fontaine, C. S. & Brown, T. A. Peat accretion histories during the past 6,000 years in marshes of the Sacramento-San Joaquin Delta, CA, USA. *Estuaries Coasts* 32, 871-892, doi: 10.1007/s12237-009-9202-8 (2009).
13. Canuel, E. A. et al. Changes in sediment and organic carbon accumulation in a highly-disturbed ecosystem: The Sacramento-San Joaquin River Delta

- (California, USA). *Marine Pollution Bulletin* 59, 154-163, doi: 10.1016/j.marpolbul.2009.03.025 (2009).
14. Brown, L. R. Will tidal wetland restoration enhance populations of native fishes? *San Francisco Estuary and Watershed Science* 1
<http://www.escholarship.org/uc/item/2cp4d8wk> (2003).
 15. Simenstad, C., Toft, J., Higgins, H., Cordell, J., Orr, M., Williams, P., Gimaldo, L., Hymanson, Z., Reed, D. Preliminary results from the Sacramento-San Joaquin Delta Breached Levee Wetland Study (BREACH). *IEP Newsletter* 12, 15-21,
<http://www.water.ca.gov/iep/newsletters/1999/1999fall.pdf> (1999).
 16. Hrodey, P. & Smith, L. The recipe for success -- what makes Liberty Island so attractive to native fish? Oral Abstracts, Bay-Delta Science Conference, September 27-29, 2010
<http://www.baydeltascienceconf.com/sciconf2010/?q=node/414> (2010).
 17. Morgan-King, T. & Schoellhamer, D. Sedimentation processes and turbidities favoring endangered fish, northern Sacramento - San Joaquin River Delta. Oral Abstracts, Bay-Delta Science Conference, September 27-29, 2010
<http://www.baydeltascienceconf.com/sciconf2010/?q=node/414> (2010).
 18. Maier, G. O. & Simenstad, C. A. The role of marsh-derived macrodetritus to the food webs of juvenile Chinook salmon in a large altered estuary. *Estuaries and Coasts* 32, 984-998, doi: 10.1007/s12237-009-9197-1 (2009).
 19. Greiner, C. M. Principles for strategic conservation and restoration. Puget Sound Nearshore Ecosystem Restoration Project Report No. 2010-01. 40 p.
http://www.pugetsoundnearshore.org/technical_papers/conservation_and_restoration_principles.pdf (2010).